

University at Albany, State University of New York

Scholars Archive

Biological Sciences

Honors College

Spring 5-2020

Body Composition of Female Collegiate Track and Field Athletes

Adelaide Cooke

University at Albany, State University of New York, acooke@albany.edu

Follow this and additional works at: https://scholarsarchive.library.albany.edu/honorscollege_biology



Part of the [Biology Commons](#), and the [Exercise Science Commons](#)

Recommended Citation

Cooke, Adelaide, "Body Composition of Female Collegiate Track and Field Athletes" (2020). *Biological Sciences*. 65.

https://scholarsarchive.library.albany.edu/honorscollege_biology/65

This Honors Thesis is brought to you for free and open access by the Honors College at Scholars Archive. It has been accepted for inclusion in Biological Sciences by an authorized administrator of Scholars Archive. For more information, please contact scholarsarchive@albany.edu.

Body Composition of Female Collegiate Track and Field Athletes

An honors thesis presented to the
Department of Anthropology,
University at Albany, State University of New York
in partial fulfillment of the requirements
for graduation with Honors in Human Biology
and
graduation from The Honors College

Adelaide Cooke

Research Mentor: Cara Ocobock, Ph.D.
Research Advisor: Adam Gordon, Ph.D.

May, 2020

Abstract

The aim of this study is to see if there is a correlation between an athlete's body composition and the demands of their different track and field events. Track and field is arguably the most diverse when it comes to body types in a single sport, with distance runners tending to be leaner and smaller and throwers more muscular and larger. However, most data collected is not female specific. Thirty-one female track and field athletes were asked to take part in a series of physical measurements, which included recording their height, weight, percent body fat, and muscle mass. They were also asked to fill out a questionnaire about their events, including person best and latest performance. This data was then used to compare muscle mass and percent fat across the event groups (distance, multis, sprints, jumps, throws) in which there was a linear relationship between the two with the exception of the multi event group. Percent muscle and percent fat were also looked at in terms of performance, in which there was no relationship. Finally, events were grouped by endurance and explosive athletes in which the endurance athletes were leaner, but the explosive athletes had more muscle mass. Across the board, there were many indicators that each event had specific body composition characteristics, but that these characteristics were not a definite indicator of performance.

Acknowledgments

Many hands have helped moved this project towards completion. First, I have to thank Dr. Ocobock for helping with its inception and then continuing to be a point of reference, even after moving to take a job at Notre Dame. Thank you to my faculty advisor Dr. Gordon for taking on this project after Dr. Ocobock left, despite not having any previous experience with working human subjects. I could not have done this without either one of you!

Thank you to the University at Albany track and field for their support, not only with this project, but also through my time as an athlete on the team. Thank you to Coach Vives for allowing me to use his athletes and thank you to the athletes that volunteered their time.

Finally, thank you to my friends and family. To the roommates that listened to me complain about the roadblocks that I encountered with this project: you mean the world to me. To my parents, thank you for advising me to take things one step at a time and for supporting me through my journey at UAlbany, not matter what it threw at me. And to Doug, my brother: never change.

List of Tables

Table 1 Average body composition values with standard deviations across event groups.....	5
--	---

List of Figures

Figure 1 Average muscle mass vs. % fat across track events	5
Figure 2 Relationship between performance and body composition in distance athletes	7
Figure 3 Relationship between performance and body composition in throwers	7

Table of Contents

Abstract	ii
Acknowledgements	iii
List of Tables	iv
List of Figures	v
Introduction	1
Methods	2
Subjects.....	2
Procudure.....	3
Results	4
Event Body Composition.....	4
Performance.....	6
Discussion	8
References	10

Introduction

Body composition of athletes has always been a topic of interest for sports medicine researchers. There are morphological and physiological characteristics that lead to a successful career in a given sport. Anecdotally, most players in the NBA are well over six-feet tall, most swimmers have broad shoulders, most football linemen are massive, and most long-distance runners are small and lean. This is largely due to the demands of the sport and the training. For example, there is evidence that runners decrease in size the further their running event is, while a swimmer's mass remains pretty constant across all distances (Gagnon et al., 2018). This could be attributed to the buoyancy of water. There is also been evidence in football players that there is a linear relationship of lean mass and fat mass seen across positions (Melvin et al., 2014). Linebackers, who value size and strength, tend to have the highest lean and fat mass values in football, while wide receivers, who must be agile to beat defenders, typically have the lowest. Due to research like this, there has been a growing belief that an athlete has to look a certain way or weigh a certain amount to succeed in their sport.

Running and track and field has come under fire for this in the last year after runner Mary Cain spoke out against Nike's Oregon Project and world-renowned coach, Alberto Salazar. As one of the best runners in the country, she joined the team out of high school. The moment she started running for them, Salazar and his all-male coaching staff insisted that she lose weight to run faster (Cain, 2019). She experienced amenorrhea, started cutting, and had suicidal thoughts, not to mention her performance suffered. Unfortunately, many male coaches, especially in running, think that training female athletes is the same as training male athletes. However, the bodies of male and female athletes are drastically different, and therefore have to be trained differently. Healthy percent fat values for fit/athletic females range from 14-24%, while in men it

is 6-17%, a difference largely due to reproductive needs (Casey, 2013). If a running coach wants their runners to be leaner so that they move less weight, they are still going to want them to take in enough food to remain healthy and reduce their risk for injury. The point here is that the percent fat where both these ideals are met is going to differ from person to person, and it certainly differs between male and female athletes.

The problem is that much of the research done on body composition has either been solely on male athletes or a combination of both sexes without evaluation of females as a separate group. For example, a study found that track and field throwers have a percent fat value of $23.6 \pm 7.8\%$, but this number was found without differentiation between the sexes (Hirsch et al., 2016). This is likely the reason for the large standard deviation.

In this study, the focus was on female collegiate athletes and their body compositions. Track and field provides a range of body types, including light, lean runners and large, muscular throwers. The goal was not only to determine the differences that develop due to different training programs but to also note the similarities between them as athletes.

Methods

Subjects: Thirty-one female NCAA Division I athletes from the University at Albany's track and field team participated in the study (Mean \pm SD; Age = 19.8 ± 1.2 yrs., Years of track participation = 8.3 ± 3.0 yrs., Height = 1.66 ± 0.07 m, Weight = 67.4 ± 16.3 kg.). In 2019-2020, the season that this study was conducted, the team won the indoor America East Conference championship for the eighth-straight year. That same week, they were also ranked the second-best team in the Northeast region by the U.S. Track & Field and Cross-Country Coaches Association behind Harvard. All athletes were over the age of 18 and signed an informed consent

approved by the university's Institutional Review Board for the protection of human subjects ($n=30$). For analysis, they were divided into five different groups: 100m-400m sprint/hurdles (sprints), 800m+ (distance), high jump/long jump (jumps), shot put/weight throw/hammer throw/discus (throws), pentathlon/heptathlon/pole vault (multis) (Table 1). These groupings were based on training groups (i.e. why pole vault was classified as multis) and demands of the event (i.e. why high jump and long jump were grouped together). Distance and throws athletes, the groups with highest number of participants ($n=11$, $n=8$ respectively) as well as the greatest difference when it comes to demands of the event, were also ranked based on performance. In the jumps group, which had a small sample size, subject FA30 was thrown out as an outlier. The average percent muscle mass for the jumps was $71.5\pm 0.3\%$ and the average percent fat was $24.7\pm 0.2\%$. FA30 had a percent muscle mass of 35.3% and percent fat of 17.6%. This was 120.7 standard deviations and 35.5 standard deviations away from both averages respectively. In such a small sample size, this kind of deviation would greatly skew the data.

Procedure: Participants came into an anthropology lab at the University at Albany for one ten-minute session. It started with a questionnaire asking for their age, years of track participation, year of college, the event(s) they participate in, and their personal best and last performance in each reported event. Their height was then measured using a portable stadiometer (SECA, Chino, CA). They were then asked to step barefoot onto a Tanita Iron Man body composition scale set to the athletic mode, grab the handles, and stand with their arms relaxed by their sides. The scale sent an undetectable electrical current through their body to determine values for weight, hydration, total percent fat and total muscle mass as well percent fat and muscle mass values for each arm, each leg, and the torso. After the collection of this data, the participant was asked for three skin fold measurements at four different sites: biceps, triceps, lower

shoulder, and at the hip. Each time, the skin was gently pinched and the thickness of this pinch was measured to the nearest millimeter using skinfold calipers. From the skin fold measurement, it was possible to calculate values for body density (females 17-19 = $1.1549 - (0.0678 \times \log \text{ of sum of average skin folds})$; females 20-29 = $1.1599 - (0.0717 \times \log \text{ of sum of average skin folds})$) and percent fat (Body Fat Percentage (%) = $(495 / \text{body density}) - 450$) (Durnin & Womeersley, 1974).

An ANOVA analysis was run to determine if there was a difference in weight, body fat, and muscle mass between the different athletic groupings, controlling for height in all cases and weight for muscle and fat mass analysis. Given the small sample size, for this analysis athletes were placed into one of two groups: endurance (distance) or explosive (sprints, throws, jumps, and multis).

Results

Event Body Composition: Using the data collected from the body composition scale, averages and standard deviations were found in each even group for both total muscle mass and percent fat. For throwers, those values were $56.7 \pm 4.5\text{kg}$ and $31.4 \pm 5.7\%$ respectively (Table 1). For distance, it was $42.3 \pm 5.0\text{kg}$ and $18.7 \pm 4.2\%$, sprints were $44.0 \pm 4.1\text{kg}$ and $20.0 \pm 2.7\%$, jumps were $50.0 \pm 4.1\text{kg}$ and $24.7 \pm 0.2\%$, and for the multis, the averages were $50.8 \pm 6.5\text{kg}$ and $21.5 \pm 2.5\%$ (Table 1).

Table 1. Average body composition values with standard deviations across event groups. Throws have the highest numbers across all categories, while distance have the smallest.

Event	n	Height (m)	Weight (kg)	% Fat	Total MM (kg)	% MM	Avg. MM Arms (kg)	Avg. MM Legs (kg)	Avg. MM Torso (kg)
Multi	4	1.69±0.09	68.2±8.0	21.5±2.5	50.8±6.5	74.5±2.4	2.7±0.4	8.9±0.9	28.1±3.8
Jumps	3	1.76±0.06	69.9±5.5	24.7±0.2	50.0±4.1	71.5±0.3	2.5±0.3	8.6±0.5	27.8±2.4
Distance	11	1.64±0.04	54.8±7.3	18.7±4.2	42.3±5.0	77.5±3.8	2.1±0.4	7.3±0.7	23.6±2.9
Throws	8	1.68±0.05	87.9±12.4	31.4±5.7	56.7±4.5	65.2±5.4	3.4±0.5	10.0±1.0	29.8±2.1
Sprints	4	1.62±0.05	58.0±4.9	20.0±2.7	44.0±4.1	75.9±2.6	2.3±0.3	7.5±0.6	24.6±2.3

These points were graphed along with the average for total muscle mass across groups (48.3kg) and the average for percent fat across all groups (22.9%). With all groups, the coefficient of determination for this relationship was $R^2 = 0.8488$ (Figure 1). However, it was obvious that the multis were an outlier. The coefficient of determination for muscle mass and percent fat without the multis was $R^2 = 0.9904$ (Figure 1). The average slope for this relationship is 0.4921.




Figure 1. The multis appears to be an outlier when it comes to the comparison between muscle mass and percent fat. There is less of a correlation when they are included in the trendline between total muscle mass and percent fat.

Unsurprisingly, the throwers were the most massive, while the distance runners were the least massive. Throwers had an average weight of 87.9 ± 12.4 kg and distance runners 54.8 ± 7.3 kg (Table 1). Throwers also tended to have more muscle mass in their arms, on average 3.4 kg compared to the 2.7 kg found with the multis, the second most massive group. While the throwers had the most muscle mass in their torso, the variability for torso muscle mass as a whole was less. Both the multis and jumps were within one standard deviation of the throwers' average of 29.8 kg. The runners were the leanest. Distance runners had an average percent fat of $18.7 \pm 4.2\%$, and sprinters had an average of $20.0 \pm 2.7\%$ (Table 1). This also meant they had the highest percentage of muscle, though not the greatest amount of muscle. There also seems to be more bilaterally symmetry when it comes to muscle mass of the legs in these two event groups compared to the other event groups. The highest percent fat values were found in the legs.

In the comparison of endurance and explosive athletes, endurance athletes had significantly higher muscle percentage ($p=0.002$). Endurance athletes had $77.5 \pm 3.9\%$ compared to the $70.4 \pm 6.2\%$ of explosive athletes. Explosive athletes also had significantly more muscle mass ($p=0.001$). They had 51.7 ± 7.0 kg compared to the 42.4 ± 5.3 kg found in endurance athletes.

Performance Based on reported performances (both personal best and most recent performance), throws and distance athletes were ranked. Distance athletes were ranked from 1-11 (1 being the best athlete). As their ranking increased, the trend was that the percent muscle mass and percent fat did as well. However, the coefficient of determination for muscle mass was $R^2 = 0.0271$ and the coefficient of determination for percent fat was $R^2 = 0.0126$ (Figure 2).




Figure 2. Percent muscle mass generally decreases and percent fat generally increase as the performance ranking increase with distance runners. However, there is not enough evidence to draw a correlation.

Throwers were ranked 1-8 (with 1 being the best). As their performance ranking increased, their percent muscle mass tended to decrease and percent fat tended to increase. However, again, the coefficients of determination for these two relationships were quite small ($R^2 = 0.0255$ and $R^2 = 0.2054$ respectively) (Figure 3).




Figure 3. Percent muscle mass generally decreases, and percent fat generally increases as performance ranking increases in throwers. However, there is not enough evidence to draw a correlation.

Discussion

The measuring and analysis of body composition of female collegiate track and field athletes in this study suggested that each event has its own morphological and physiological characteristics. However, body composition did not factor into performance in a given event. The body composition trends were similar to the findings of other studies done on NCAA Division I track and field athletes, such as Hirsch et al. (2016). The throwers have the greatest muscle mass and percent fat. This makes sense from a physics standpoint. If the aim is to generate as much power, you want to create as much kinetic energy as possible before you throw (Bartoniets, 1994). An increase in mass or velocity increases kinetic energy. The trick of throwing is transferring this kinetic energy into the throwing implement. This is why the successful smaller throwers must be quicker than their competitors. It should also be noted that though the correlation was far from strong when it came to performance and body composition in throwers, the percent fat decreases with increased performance ranking. This suggests that an increase in mass to increase performance is not about increasing body fat but increasing lean body mass.

On the other end of the spectrum are the runners. As Gagnon et al. (2018) proved, as you increase the running distance you are training for, you decrease your mass. This is true with the sprinter and distance groups, with distance being the less massive of the two. This could come back to the types of muscle fibers required to be successful in each run. Slow-twitch fibers or type I muscle fibers are smaller in diameters compared to fast-twitch or type II muscle fibers (Kenney et al., 2015). Distance would have more type I fibers than sprinters, suggesting that their muscle mass would be less if a cross section had the same number of muscle fibers as a sprinter. A sprinter also has more muscle mass in their arms compared to distance (2.3 ± 0.3 kg. vs 2.1 ± 0.4 kg) (Table 1). This is because their upper body is used to balance their strides (Mann &

Sprague, 1983). They can energetically afford to have the increased weight in their upper body because they run for a much shorter time than distance runners.

This is furthered by the comparison of endurance and explosive athletes, which not only compared distance runners to the sprinters but also to other events utilizing these fast-twitch, type II muscle fibers. Though distance, endurance runners were significantly leaner, explosive athletes had significantly more muscle mass. If we directed this back to what we were talking about above, a point could be made about the different distribution of fiber-type that can be found in endurance vs. explosive athletes. Distance would be the group Mary Cain would find herself in. Despite what Coach Salazar believed, we did not find a great enough correlation between performance and percent fat or performance and muscle mass to come to the conclusion that either body composition variable contributes to an athlete's performance.

The multis appear to be an outlier in the relationship between muscle mass and percent fat. They had low fat percentages for the given muscle mass that they had. This could be due to the nature of the event they train for. A multi event challenge an athlete across several disciplines. They have to be explosive in the jumps, massive and strong in the throws, while still lean enough to run hurdles and the 800m. However, them being an outlier could also be attributed to the small sample size, which is not large enough to draw a significant conclusion.

These trends beg the question if body composition is a result of training or if individuals are drawn to certain events because of their genetically pre-disposed body type. At the University at Albany, the throwers are given one lifting program, the distance group another, while the multis, jumpers, and sprinters lift together. However, there is both diversity in the throws group and with the multis, jumpers, and sprinters. This suggests that while strength and conditioning is a vital component to building one's body composition for an event, it is likely the

individual training done with coaches that plays more of a role in performance. Furthermore, this study reiterates the importance of having female-specific studies in the field of sports medicine, which can only help coaches better train their female athletes in a healthy and successful manner.

References

- Bartonietz, K. E. (1994). Training of technique and specific power in throwing events. *Modern Athlete and Coach*, 32(1), 1-16.
- Cain, M. (2019, November 7). I was the fastest girl in America, until I joined Nike. *New York Times*. Retrieved from <https://www.nytimes.com/2019/11/07/opinion/nike-running-mary-cain.html>
- Casey, J. (2003). Body fat measurement: Percentage vs. body mass. *WebMD*. Retrieved from <https://www.webmd.com/diet/features/body-fat-measurement#2>
- Durnin, J. V., & Womersley, J. V. G. A. (1974). Body fat assessed from total body density and its estimation from skinfold thickness: measurements on 481 men and women aged from 16 to 72 years. *British Journal of Nutrition*, 32(1), 77-97.
- Fleck, S. J. (1983). Body composition of elite American athletes. *The American Journal of Sports Medicine*, 11(6), 398-403.
- Gagnon, C. M., Steiper, M. E., & Pontzer, H. (2018). Elite swimmers do not exhibit a body mass index trade-off across a wide range of event distances. *Proceedings of the Royal Society B*, 285(1882). <http://doi.org/10.1098/rspb.2018.0684>
- Hirsch, K. R., Smith-Ryan, A. E., Trexler, E. T., & Roelofs, E. J. (2016). Body composition and muscle characteristics of Division 1 track and field athletes. *The Journal of Strength & Conditioning Research*, 30(5), 1231-1238. doi: 10.1519/JSC.0000000000001203

- Kenney, W. L., Wilmore, J. H., & Costill, D. L. (2015). Structure and function of exercising muscle, in *Physiology of Sport and Exercise* (pp. 27-49). Human Kinetics.
- Mann, R., & Sprague, P. (1983). Kinetics of sprinting. *Track and Field Quarterly Review*, 83(2), 4-9.
- Melvin, N. M., Smith-Ryan, A. E., Wingfield, H. L., Ryan, E. Ed., Trexler, E. T., & Roelofs, E. J. (2014). Muscle characteristics and body composition of NCAA division I football players. *The Journal of Strength and Conditioning Research*, 28(12), 3320-3329.