

University at Albany, State University of New York

Scholars Archive

Geology Theses and Dissertations

Atmospheric and Environmental Sciences

1976

Petrology of the Oceanographer Fracture Zone (35°N35°W)

Tsugio Shibata

University at Albany, State University of New York

Follow this and additional works at: https://scholarsarchive.library.albany.edu/cas_daes_geology_etd



Part of the [Geochemistry Commons](#), [Geology Commons](#), [Oceanography Commons](#), and the [Tectonics and Structure Commons](#)

Recommended Citation

Shibata, Tsugio, "Petrology of the Oceanographer Fracture Zone (35°N35°W)" (1976). *Geology Theses and Dissertations*. 104.

https://scholarsarchive.library.albany.edu/cas_daes_geology_etd/104

This Dissertation is brought to you for free and open access by the Atmospheric and Environmental Sciences at Scholars Archive. It has been accepted for inclusion in Geology Theses and Dissertations by an authorized administrator of Scholars Archive. For more information, please contact scholarsarchive@albany.edu.

PETROLOGY OF THE OCEANOGRAPHER FRACTURE ZONE (35°N35°W)

by

Tsugio Shibata

Abstract of a Dissertation

Submitted to the State University of New York at Albany

in Partial Fulfillment of

the Requirements for the Degree of

Doctor of Philosophy

College of Arts and Sciences

Department of Geological Sciences

1976

ABSTRACT

During a geological and geophysical survey of the Oceanographer Fracture Zone (35°N, 35°W), seventeen dredge hauls containing a variety of rocks were obtained. Petrographic study shows that these rock samples can be classified into six main rock types: fresh basalt, weathered basalt, metabasalt, gabbro, metagabbro, and serpentinite. Most of the dredge hauls were positioned on the steep, southern wall of the fracture zone, and an inference from the dredging results suggests that basalt is the most abundant rock type which outcrops at the junction between the rift valley and the fracture zone; however, as we move away from the junction, metabasalts and metagabbros apparently become more abundant in the outcrops on the wall of the fracture zone.

Fresh basalts from the Oceanographer Fracture Zone are characterized by the petrography, mineralogy, and major element chemistry typical of abyssal tholeiites, but are somewhat enriched in large ion lithophile elements (K, La, etc.). The major constituent minerals observed in these fresh tholeiitic basalts are olivine, plagioclase, clinopyroxene, and opaque minerals. Compositional variation of olivine within a given specimen is small, and fayalite contents steadily increase as FeO^* (total iron as FeO)/MgO in the host basalt increases. Clinopyroxenes of these tholeiitic basalts are mostly augites, and no Ca-poor pyroxene is observed. Plagioclase including those that occur in the groundmass show an overall compositional range from An_{86} to An_{44} . There is a positive correlation between albite content and $\text{Fe}/(\text{Fe} + \text{Mg})$ in these plagioclases; i.e., $\text{Fe}/(\text{Fe} + \text{Mg})$ increases as plagioclase becomes more sodic.

Major element analyses of tholeiitic basalts from two adjacent

dredge hauls (RD 7 and 8) show that these tholeiitic basalts are closely related chemically and that they apparently represent points on a continuous liquid line of descent. This compositional variation can be ascribed to fractional crystallization of phenocryst phases observed in these tholeiites.

Metamorphic rocks from the Oceanographer Fracture Zone range in metamorphic grade from zeolite to greenschist facies. Metabasalts are massive and show no evidence of deformation; metagabbros are to some degree deformed. Petrographic study of these metagabbros indicates that plagioclase grains are commonly strained, bent, and ruptured, whereas pyroxene and hornblende grains are replaced by a metamorphic mineral assemblage made up mainly of actinolite and chlorite. Apparently, plagioclase responded differently than the mafic minerals to these tectonic and metamorphic conditions.

The Oceanographer rock collection is characterized by abundant rocks with the hornblende plus plagioclase assemblage. It is suggested that this rock type must be considered as one of the potential candidates for the constitution of oceanic crust.

PETROLOGY OF THE OCEANOGRAPHER FRACTURE ZONE (35°N35°W)

by

Tsugio Shibata

A Dissertation

Submitted to the State University of New York at Albany

in Partial Fulfillment of

the Requirements for the Degree of

Doctor of Philosophy

College of Arts and Sciences

Department of Geological Sciences

1976

TABLE OF CONTENTS

	Page
ABSTRACT	1
LIST OF FIGURES	iv
LIST OF PLATES	v
LIST OF TABLES	vi
ACKNOWLEDGMENTS	vii
CHAPTER I	
INTRODUCTION	1
CHAPTER II	
REGIONAL SETTING AND DREDGE HAULS	5
Contents of Dredge Hauls and Rock Association on the Southern Wall	7
CHAPTER III	
FRESH BASALTS	15
Petrography	15
RD6	15
RD7 and 8	15
RD10	17
Mineralogy	19
Olivine	20
Pyroxene	25
Plagioclase	35
Chemistry	41
Fractionation of the Oceanographer Tholeiite Magma	48
Discussion	58
CHAPTER IV	
METAMORPHOSED BASALTS AND GABBROS	64
Petrography	64
Metabasalts	64
Metadolerites	65
Metagabbros	66
Metabasites	67
Cataclastic Textures of Metagabbros	69
Discussion	76
CHAPTER V	
SUMMARY AND CONCLUSION	79

	Page
BIBLIOGRAPHY	82
APPENDIX I Phenocryst - Bulk Rock Composition Relations of Abyssal Tholeiites and Their Petrogenetic Significance	88
II Electron Microprobe Analyses of Olivine, Plagioclase and Clinopyroxene	<i>in pocket (27pp)</i>

LIST OF FIGURES

Figure Number	Page
1. Index Map Showing the Location of the Oceanographer Fracture Zone	6
2. Generalized Contour Map of the Oceanographer Fracture Zone Showing the Location of the Seventeen Dredge Hauls	10
3. Dredge Depths and Recovered Rock Types	11
4. Frequency Diagrams Showing the Compositional Range of Olivines Observed as Phenocrysts or Microphenocrysts in the Oceanographer Tholeiites	23
5. Pyroxene Quadrilateral Showing the Plots of Clinopyroxenes from the Oceanographer Tholeiites	28
6. Relation Between Si and Al in Clinopyroxenes from the Oceanographer Tholeiites	31
7. Relation Between Al and Ti in Clinopyroxenes from the Oceanographer Tholeiites	32
8. Ol-Di-Qz Diagram Showing Plots in Molecular Proportions of Clinopyroxenes	33
9. Di-01-Pl Diagram Showing Plots in Molecular Proportions of Clinopyroxenes	34
10. A: The Sum of Fe and Mg in Plagioclases Plotted Against Anorthite Contents	39
B: The Ratio of Fe to Fe plus Mg in Plagioclases Plotted Against Anorthite Contents	39
11. Potassium Contents in Plagioclases	40
12. AFM Diagram $[(Na_2O + K_2O) : FeO^* : MgO]$ Showing the Distribution of the 48 Analyses of RD 7 and 8 Tholeiites	49
13. Oxide Percents Plotted Against the Solidification Index	50
14. AFM Diagram $[(Na_2O + K_2O) : FeO^* : MgO]$ Comparing the Fractionation Trend Defined by the Oceanographer Samples from Dredges 7 and 8 With Four Other Synoptic Analyses of Dredge Populations	60
15. AFM Diagram $[(Na_2O + K_2O) : FeO^* : MgO]$ Comparing the Fractionation Trend Defined by the Oceanographer Samples from Dredges 7 and 8 with Four Well-Studied Tholeiitic Basalt Suites	62

LIST OF PLATES

Plate Number	Page
1. Porphyritic Basalt	16
2. Photomicrograph Showing the Glass-Rich Crust of Basalt Fragment	18
3. Microstructure of Plagioclase Mylonite	68
4. Metamorphosed Olivine Gabbro	71
5. Recrystallization Along Plagioclase Grain Boundaries	72
6. Fractured Plagioclases in Metagabbro	73
7. Metagabbro with Angular Plagioclase Fragments	74

LIST OF TABLES

Table Number	Page
1. Location of Dredge Stations and Recovered Rock Type	8
2. Selected Analyses of Olivine	21
3. Average Olivine Compositions and FeO/MgO of the Host Rocks	24
4. Selected Analyses of Pyroxene	26
5. Selected Analyses of Plagioclase	36
6. Chemical Analyses and Norms of Six Abyssal Tholeiites	42
7. Atomic Absorption Analyses for the Oceanographer Tholeiites	44
8. Fractionation Models	53

ACKNOWLEDGMENTS

I am grateful to A. Miyashiro for suggesting this thesis and for providing support, advice, and encouragement throughout my graduate studies. P. J. Fox and S. E. DeLong also offered much-needed encouragement and motivation during the course of this study. A. Miyashiro, S. E. DeLong, and P. J. Fox are gratefully acknowledged for critically reading the manuscript and making many suggestions which helped to improve its quality in many ways. I also wish to acknowledge Captain H. Kohler, the officers and crew of the R/V VEMA for aiding in the collection of rock samples. F. Shido kindly assisted with the chemical analyses of rock specimens and S. E. DeLong with the microprobe analyses of minerals. J. A. Karson provided me with excellent assistance and companionship during my shipboard work.

This study was financially supported by National Science Foundation Grants A020742-001 and OCE75-21753.